

<https://helda.helsinki.fi>

Partial restoration of normal intestinal microbiota in morbidly obese women six months after bariatric surgery

Koffert, Jukka

2020-11-26

Koffert , J , Lahti , L , Nylund , L , Salminen , S , Hannukainen , J C , Salminen , P , de Vos , W M & Nuutila , P 2020 , ' Partial restoration of normal intestinal microbiota in morbidly obese women six months after bariatric surgery ' , PeerJ , vol. 8 , 10442 . <https://doi.org/10.7717/peerj.10442>

<http://hdl.handle.net/10138/322895>

<https://doi.org/10.7717/peerj.10442>

cc_by

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.



Partial restoration of normal intestinal microbiota in morbidly obese women six months after bariatric surgery

Jukka Koffert^{1,2,*}, Leo Lahti^{3,*}, Lotta Nylund^{4,5}, Seppo Salminen⁴, Jarna C. Hannukainen², Paulina Salminen⁶, Willem M. de Vos^{7,8} and Pirjo Nuutila^{2,9}

¹ Department of Gastroenterology, Turku University Hospital, Turku, Finland

² Turku PET Centre, University of Turku, Turku, Finland

³ Department of Mathematics and Statistics, University of Turku, Turku, Finland

⁴ Functional Foods Forum, University of Turku, Turku, Finland

⁵ Food Chemistry and Food Development, Department of Biochemistry, University of Turku, Turku, Finland

⁶ Division of Digestive Surgery and Urology, Turku University Hospital, Turku, Finland

⁷ RPU Immunobiology and Human Microbiome, University of Helsinki, Helsinki, Finland

⁸ Laboratory of Microbiology, Wageningen University and Research, Wageningen, The Netherlands

⁹ Department of Endocrinology, Turku University hospital, Turku, Finland

* These authors contributed equally to this work.

ABSTRACT

We studied the impact of bariatric surgery on the intestinal microbiota of morbidly obese study subjects. A total of 13 morbidly obese women (five of which had type 2 diabetes) and 14 healthy age- and gender-matched controls were recruited and the microbiota composition of fecal samples were determined by using a phylogenetic microarray. Sampling of the patients took place just one month before and 6 months after the operation. Within six months after bariatric surgery, the obese subjects had lost on average a quarter of their weight whereas four of the five of the diabetic subjects were in remission. Bariatric surgery was associated with an increased microbial community richness and Bacteroidetes/Firmicutes ratio. In addition, we observed an increased relative abundance of facultative anaerobes, such as *Streptococcus* spp., and a reduction in specific butyrate-producing Firmicutes. The observed postoperative alterations in intestinal microbiota reflect adaptation to the changing conditions in the gastrointestinal tract, such as energy restriction and the inability to process fiber-rich foods after bariatric surgery.

Subjects Microbiology, Clinical Trials, Diabetes and Endocrinology, Nutrition, Metabolic Sciences

Keywords Fecal microbiota, Bariatric surgery, Type 2 diabetes, Obesity, Clinical trial

INTRODUCTION

The global obesity epidemic is still worsening affecting a total of 603.7 million adults and 107.7 million children with ominous implications for both physical and economic health (*Collaborators GBDO et al., 2017*). Several studies have shown that intestinal microbes contribute to the regulation of energy homeostasis, fat storage, and energy uptake, and are likely to play a role in obesity and its complications (*Karlsson et al., 2013; Kootte et al.,*

Submitted 29 May 2020

Accepted 6 November 2020

Published 26 November 2020

Corresponding author

Pirjo Nuutila, pirjo.nuutila@utu.fi

Academic editor

Stefano Menini

Additional Information and
Declarations can be found on
page 12

DOI 10.7717/peerj.10442

© Copyright
2020 Koffert et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

2012; *Krajmalnik-Brown et al., 2012*). Indeed, the intestinal microbiota signatures, such as altered composition as well as reduction in taxonomic diversity and functional richness have been associated with obesity (*Cotillard et al., 2013; Turnbaugh et al., 2009*) and several experimental studies have shown a causal link between obesity and gut microbiota (*Liou et al., 2013; Ridaura et al., 2013*). In addition, aberrations of intestinal microbiota have been shown to be associated with the serum metabolome and may contribute to insulin resistance in non-diabetic subjects (*Pedersen et al., 2016*).

Initial studies suggested that obesity could be associated with an increased relative abundance of Firmicutes and decreased levels of Bacteroidetes (*Furet et al., 2010; Ley et al., 2006*). However, more recent studies have reported results ranged from an increase in total diversity to absence of change and even decrease in alpha-diversity (*Damms-Machado et al., 2015; Louis et al., 2016; Patil et al., 2012*). These may be explained by differences in diet, host genotype and physiology and the presence of comorbidities, such as type 2 diabetes (T2D) (*Remely et al., 2016*). Moreover, these studies addressed the number and not the activity of these bacterial phyla. A more recent study addressing the fecal metaproteome showed that bacteria belonging to the Bacteroidetes were less abundant, but metabolically much more active in morbidly obese versus non-obese subjects (*Kolmeder et al., 2015*).

In patients with morbid obesity bariatric surgery is currently the only effective treatment with long-term results of sustainable weight loss and remission or amelioration of obesity-related comorbidities (*Adams et al., 2017; Puzziferri et al., 2014; Schauer et al., 2017; Sjostrom, 2013*). The underlying mechanisms behind the surgically induced weight loss have not been fully elucidated. However, reduced food ingestion, increased satiety, increased gastric emptying and shift in bile acid metabolism have been suggested to have a role (*LaFerrere et al., 2008; Plum et al., 2011*). In addition, increased diversity and changes in community composition of the intestinal microbiota have been reported after bariatric surgery, thus suggesting that some of its beneficial effects may be mediated by intestinal microbiota (*Damms-Machado et al., 2015; Graessler et al., 2014; Palleja et al., 2016*). Previously published studies on the effect of bariatric surgery on fecal microbiota are not consistent and suggest contradictory results concerning changes in fecal microbiota diversity after the surgery and between different surgery types (*Lee et al., 2019; Paganelli et al., 2019; Sanchez-Alcoholado et al., 2019*). The objective of this study was to characterize the microbiota of morbidly obese women and to evaluate the impact of bariatric surgery on microbiota composition.

STUDY SUBJECTS AND METHODS

Study population

Morbidly obese study subjects were recruited from patients undergoing bariatric surgery procedure as part of their normal treatment at the Hospital District of Southwest Finland (*Dadson et al., 2017; Koffert et al., 2018*). The surgical inclusion criteria were age between 18–60 years and BMI ≥ 40 kg/m² (or ≥ 35 kg/m² with an additional obesity-related comorbidity). In addition to standard exclusion criteria for bariatric surgery patients with T2D requiring insulin treatment were excluded. Healthy study subjects were recruited to




Figure 1 Study flowchart. (A) Healthy study subjects. (B). Obese study subjects.

[Full-size DOI: 10.7717/peerj.10442/fig-1](https://doi.org/10.7717/peerj.10442/fig-1)

study protocol by advertisements in local newspaper from January to December in 2011. The inclusion criteria for the control population were body-mass index 18–27 kg m², age 18–60 years, fasting plasma glucose less than 6.1 mM, and normal oral glucose tolerance test. Altogether 42 females were recruited for the study, including 27 obese subjects and 15 controls. For nine obese subjects, one of the time points was discarded due to the inability to PCR amplify the extracted DNA or low reproducibility of the phylogenetic microarray results (see below). Moreover, three obese subjects withdrew from the study for personal reasons, and further two subjects who had a successful microbiota profile only from a single time point were discarded. This yielded the final data set with 13 obese subjects who had successful microbiota profiles from both time points and 14 controls that were included in the analysis (Fig. 1). Of the 13 obese subjects, five had T2D and in eight had normal glycaemia. Only three study subjects with type 2 diabetes were on metformin medication. Written informed consents were obtained prior the studies. The protocol (NCT01373892/SLEEVEPET) was approved by the Ethics Committee of the Hospital District of Southwest Finland (ETMK 99/180/2010). The clinical study was performed at the Turku PET Centre, the University of Turku and Turku University Hospital (Turku, Finland) between January 2011 and October 2013 and fecal microbiota samples analyzed in 2014.

Study design

At screening visit, study subjects gave their signed approval prior to their enrolment in to the study. The medical history was recorded, the physical examination, anthropometric measurements, blood and oral glucose tolerance tests were performed, and instructions for fecal sample collection were provided. On a separate study visit abdominal fat masses

were assessed with MRI. Thereafter, obese subjects followed a very low-calorie diet (VLCD; 800kcal/day) for one month before the bariatric operation. Of the morbidly obese subjects, seven underwent sleeve gastrectomy and six Roux-en-Y gastric bypass (RYGB) surgery. All the morbidly obese study subjects received perioperative antibiotic Cefuroxime 3g intravenously. The post-operational phase was conducted six months after the surgery and all studies were repeated as in baseline. Food diaries were used to document dietary intake in healthy controls and before the VLCD and 6 months postoperatively in obese subject.

Fecal sample collection and microbiota profiling

Fecal samples for the controls were collected in a single time point and for the obese subjects both before the VLCD (one month before the surgery) and six months after the surgery. DNA from fecal samples was extracted with repeated bead-beating as described previously (Nylund *et al.*, 2010) and used for phylogenetic profiling with the Human Intestinal Tract (HIT) Chip, a custom-made Agilent phylogenetic microarray (Rajilic-Stojanovic *et al.*, 2009). The HITChip analysis allows phylotype-level profiling at a resolution of ~0.1% relative abundance with >98% reproducibility (Claesson *et al.*, 2009) based on >5,000 probes designed to target 16S rRNA gene sequences of over 1,000 intestinal bacterial phylotypes from 130 genus-like groups (defined by >90% 16S rRNA gene similarity). Further details on the sample processing, microarray, and hybridization protocols were reported previously (Rajilic-Stojanovic *et al.*, 2009). The microarray hybridization data was extracted from scanned microarray images with the Agilent Feature Extraction software. The technical replicates were quantile-normalized with custom R scripts as previously described (Lahti *et al.*, 2014), and reproducible hybridizations (Pearson $r > 98\%$) were accepted for the analysis and stored in a custom MySQL database. Between-sample normalization was performed at probe level with the minimum–maximum method, and probe summarization at the phylotype-level (>98% 16S rRNA sequence similarity) was done with the frozen Robust Probabilistic Averaging (fRPA) method (Lahti *et al.*, 2011; Lahti *et al.*, 2013) and then to the genus (>90% 16S rRNA sequence similarity) and order-like 16S rRNA gene sequence groups by summing up phylotype abundances as described earlier (Lahti *et al.*, 2014). The processing scripts are available at <https://github.com/microbiome/HITChipDB>. In this work, we focus primarily on genus-level variation.

Statistical analysis

The HITChip profiles were centered log-ratio transformation (CLR)-transformed before statistical testing to remove compositionality bias (Aitchison, 1986), and Euclidean distances for the CLR-transformed genus-level abundances were used as the beta diversity measure. For hierarchical clustering, we used complete linkage with Spearman correlation. The surgery effect on individual taxa between the matched obese subject time points 1–2 was quantified with a random effects linear model with subject as the random effect and time as the fixed effect (lmer R package). Unpaired *t*-test was used for two-group comparisons. Multi-group comparisons were performed with ANOVA and the Tukey post-hoc test. In order to reduce multiple testing, we removed from comparisons the rare taxa that were

observed with <10% prevalence at >0.1% relative abundance. To further support the analysis, we verified the multi-group comparisons with non-parametric Kruskal-Wallis test and diagnostic plots. PERMANOVA (adonis in the *vegan* R package) was used for the group-wise community comparisons. The *p*-values were adjusted for multiple corrections with Benjamini–Hochberg method. The FDR<0.05 was considered statistically significant, and FDR<0.25 borderline significant. Spearman correlation of the probe-level HITChip signals were used for subject identification based on best match between the two timepoints. Microbiota richness and diversity were quantified from the probe-level HITChip data by observed probe count and Shannon index, respectively. Tools from the phyloseq (McMurdie & S, 2013) and microbiome (<https://microbiome.github.io/tutorials/>) R packages were used to support analysis and visualization.

Sample size for original study protocol in PET-imaging was determined by joint *a priori* power analysis based on longitudinal PET studies using 18F-FTHA (Karmi et al., 2010; Viljanen et al., 2009) which suggested that a sample size of 10+10 would be sufficient for establishing the predicted effects at *P*<0.05 with actual power exceeding 0.95. These study subjects were used for intestinal microbiota analysis.

RESULTS

Effect of bariatric surgery on glucose metabolism

Before surgery, the obese study group was insulin resistant and had increased systemic inflammation compared to healthy controls (Table 1). In addition, the obese study subjects had higher HbA1c and 2-hour glucose concentrations in OGTT compared to healthy controls (Table 1). Bariatric surgery resulted in marked weight loss (on average 23%) with improved insulin resistance (Table 1). Furthermore, diabetes remission occurred in 80% (4/5) of the T2D subjects after surgery.

Dietary intake preceding and after surgery

Mean caloric intake was lower in obese subjects compared to healthy controls but the difference was only marginally significant ($1,304 \pm 570$ kcal vs. $1,625 \pm 518$, *p* = 0.07, respectively). The total calorie intake decreased in obese subject by 35% on average after bariatric surgery (*p* < 0.01). Nutrient intake was lower postoperatively, while carbohydrate-, protein- and fat-intake decreased on average by 27% (*p* < 0.01), 18% (*p* < 0.01) and 6% (*p* = 0.04) respectively.

Control subjects form two distinct groups

Unsupervised hierarchical clustering of the genus-level profiles indicated two distinct subgroups among the control subjects (Fig. S1). These included 4 normal-weight control subjects with mean BMI 24.9 kg/m^2 (group A) and 10 “lean” control subjects with a lower average BMI of 21.5 kg/m^2 (group B). Differences in body fat and energy consumption were not significant. The group A controls had on average higher levels of bacteria related to *Allistipes*. In contrast, the group B controls had on average higher levels of bacteria belonging to *Ruminococcus obeum*, *Coprococcus eutactus* and *Clostridium nexile* (*p* < 0.25).

Table 1 Data are presented as mean \pm SD.

Anthropometrics	Controls (<i>n</i> = 14)	Obese (<i>n</i> = 13)		
		Presurgery	<i>p</i> -value	Postsurgery
Sex (female/male)	14/0	13/0		
Age (years)	44 \pm 10	43 \pm 11	0,88	44 \pm 10
Weight (kg)	62 \pm 7.3	113 \pm 17*	<0.01	86 \pm 14
BMI (kg m ⁻²)	22.8 \pm 2.9	40.9 \pm 4.2*	<0.01	31.4 \pm 3.6*
Body fat (%)	32 \pm 5.8	49 \pm 5.4*	<0.01	41 \pm 3.7
FFA (mM)	0.50 \pm 0.20	0.73 \pm 0.29*	0.62	0.78 \pm 0.24*
Fasting glucose (mM)	5.3 \pm 0.6	6.0 \pm 1.1	0.11	5.4 \pm 0.9
Fasting insulin (mU L ⁻¹)	5.6 \pm 3.6	11.0 \pm 7.9*	0.19	7.6 \pm 5.0
OGTT 2 h glucose (mM)	5.6 \pm 1.2	8.7 \pm 2.8*	<0.01	4.8 \pm 2.0
T2D (<i>n</i>)	0	5		1
uCRP (mg*l ⁻¹)	0.8 \pm 1.0	2.9 \pm 2.5*	0.013	0.9 \pm 1.0
HbA1c (%)	5.6 \pm 0.3	5.8 \pm 0.7	0.01	5.3 \pm 0.5
HOMAIR (fraction)	1.1 \pm 0.9	3.1 \pm 2.7*	0.18	1.9 \pm 1.5

Notes.

HOMAIR, homeostatic model assessment for insulin resistance; FFA, free fatty acids; OGTT, oral glucose tolerance test; T2D, type 2 diabetes.

**p* < 0.05 for obese patients pre- and postintervention when compared to controls in Student's *t*-test.

Surgery type has no effect on microbiota composition

No differences were observed between the alternative surgery procedures (bypass *n* = 6; sleeve *n* = 7) in post-surgery samples in community composition quantified by the overall beta diversity or individual genus-like groups (Fig. S2A). These groups were combined in the subsequent microbiota analyses.

Microbial richness shifts towards controls in patients following surgery

Before surgery, the average richness was significantly lower in the obese subjects compared to the group A controls (*p* = 0.01) and borderline significant compared to the group B controls (*p* = 0.12) (Fig. 2). After the surgery, the difference to group A controls was borderline significant (*p* = 0.14), and difference to the group B controls was not significant (*p* = 0.87).

Microbiota community structure shows changes following bariatric surgery

Bariatric surgery was associated with a shift in the Bacteroidetes-Firmicutes ratio towards that of the control subjects (Fig. 2B). The overall differences in microbiota composition (beta diversity) between the obese and control subjects were not significant after the surgery. Notably, whereas the average abundance of Firmicutes decreased after surgery, specific members of the Firmicutes exhibited qualitatively different increasing shifts following the surgery (Fig. 3), indicating that the phylum-level comparison of the Bacteroidetes-Firmicutes ratio should be complemented by the analysis of specific members within these groups.




Figure 2 Microbiota richness in the obese and control groups. (A) The group-wise differences are significant for richness ($p = 0.01$), and borderline significant for diversity ($p = 0.22$). At the baseline time point (before surgery), the richness was lower in the obese subjects compared to the normal-weight controls ($p = 0.01$) and lean controls (borderline significant; $p = 0.12$). After the surgery, microbiota richness in the obese reached similar levels with the lean controls ($p = 0.87$) but remained somewhat lower compared to the normal-weight controls ($p = 0.14$). (B) The surgery was associated with a significant shift in the Bacteroidetes-Firmicutes ratio ($p = 0.04$), whereas the other differences were not significant.

Full-size [DOI: 10.7717/peerj.10442/fig-2](https://doi.org/10.7717/peerj.10442/fig-2)

Genus-level shifts following the surgery

At the level of specific bacterial genera, bariatric surgery was associated with an increased abundance of specific members of bacteria belonging to the Bacteroidetes, *Streptococcus* spp. (*S. mitis* and *S. bovis*), as well as *Sporobacter termitidis* (*Clostridium* cluster IV) (Fig. 4), so that the abundances of these bacteria were shifting towards the group A controls. The observed shift towards these controls was evident also as a reduced abundance in specific members of the *Clostridium* cluster XIVa (including bacteria related to *Ruminococcus obeum*, *Coprococcus eutactus*, and other *Clostridium* groups) after surgery.

Individuality of microbiota composition

Microbial composition remained individual-specific and identifiable following surgery for most (11/14; 79%) subjects based on the closest match with the pre-surgery samples (see Methods). However, three subjects experienced so remarkable changes during the intervention that reliable subject identification based on microbiota composition was not possible after the surgery (best match was with a sample from another subject; Fig. S3). However, for two of these incorrectly identified subjects, the second best match was with the correct subject. For the third misidentified obese subject, his/her own pre-surgery sample was only the 9th best match, suggesting drastically altered microbiota composition or potential shift in community type following the surgery.




Figure 3 Community similarity before and after bariatric. Sample ordination by genus-level community composition illustrates a significant shift from obese microbiota (red) towards group A control group (dark blue) following surgery ($p < 0.05$). The lower panel shows the ten most strongly associated genus-like groups for each ordination axis, indicating that surgery is, on average, associated with a reduction of genera from the *Clostridium* cluster XIVa and Proteobacteria groups, and an increase in Bacteroidetes, *Clostridium* cluster IV, and uncultured Clostridiales. For targeted analysis of changes in genus-level groups following surgery, see Fig. S2B.

Full-size [DOI: 10.7717/peerj.10442/fig-3](https://doi.org/10.7717/peerj.10442/fig-3)

Effect of diabetes status

In addition, the microbiota characteristics of metformin-treated obese subjects ($n = 3$) and obese subjects with no metformin treatment ($n = 10$) were compared at the baseline time point. Diabetes status did not have a statistically significant effect on the overall microbiota composition as quantified by the standard permutation analysis of variance (PERMANOVA; $p = 0.28$). However, metformin was associated with a significant reduction in intestinal microbiota diversity ($p = 0.02$; Fig. 5). A similar non-significant trend is seen in microbiota richness ($p = 0.21$).

DISCUSSION

In this study, the intestinal microbiota composition of morbidly obese females with and without T2D and the effects of bariatric surgery induced weight loss on intestinal microbiota composition and diversity were evaluated. The effect of bariatric procedure on microbiota




Figure 4 Genus-level microbiota signatures among the study subjects. Abundance variation in the genus-level groups that exhibited significant changes following the surgery. Z-transformed \log_{10} relative abundances (HITChip signal) are shown for each genus. The color code indicates increased (red) or decreased (blue) abundance compared to the overall mean abundance of the genus across all samples.

[Full-size](#) DOI: 10.7717/peerj.10442/fig-4




Figure 5 Metformin treatment and microbiota diversity. Metformin treatment is associated with reduced microbiota diversity ($p = 0.02$).

[Full-size](#) DOI: 10.7717/peerj.10442/fig-5

signatures was characterized by increased microbiota richness, increased abundance of facultative anaerobes and decreased abundances of specific butyrate-producing Firmicutes.

Before bariatric surgery, the overall microbiota composition of the morbidly obese subjects was closer to the group B controls than group A controls. This potentially reflects the effect of calorie-restricted weight-loss diet in the obese group (Cotillard et al., 2013) as

the patients undergoing bariatric surgery in Finland are required to lose approximately 5% of their weight before surgery. Six months after the operation, microbiota composition of the obese subjects had shifted towards the group A controls (Fig. 3). These shifts were observed both in the overall community composition (beta diversity), in specific genera, community richness and in Bacteroidetes-Firmicutes ratio. Despite the shifts observed, microbial composition remained individual-specific and identifiable following surgery for the most which is in line with current literature (Durack & Lynch, 2019; Healey et al., 2017). In a recent study conducted by Paganelli and her colleagues low-calorie diet and bariatric surgery per se only temporarily declined fecal microbiota diversity and gradually these changes returned to baseline levels in six months (Paganelli et al., 2019). On the other hand, Lee's research group documented increase in fecal microbiota diversity after the RYGB and drug-induced weight loss (Lee et al., 2019). All of these studies are limited by low number of study subjects and shortage of male study subjects in different intervention groups.

In previous studies, subjects with prediabetes or overt T2D have been reported to have a distinct microbiota profile compared to non-diabetic subjects (Allin et al., 2018; Qin et al., 2012). In current study, we did not observe differences in microbiota signatures related to the diabetic status at the baseline ($p = 0.048$). However, microbiota diversity was significantly decreased in metformin—treated diabetic subjects compared to diabetic subjects without medication (Fig. 5). This finding is in contrast to recent study by Forslund and colleagues, where decreased bacterial richness was detected among metformin-naïve individuals with type 2 diabetes when compared to individuals treated with metformin (Forslund et al., 2015).

Increased relative abundances of facultative anaerobic microbes, such as *Streptococcus* spp. and *Allistipes* et rel. were observed after bariatric surgery. This is in line with previous studies reporting increased relative abundances of facultative anaerobic microbes from upper gastrointestinal tract (such as *Streptococcus*, *Veillonella* and *Escherichia* spp.) after bariatric surgery (Ilhan et al., 2017; Pallega et al., 2016). Paganelli's research group showed that after the VLCD relative abundance of Streptococcae was decreased but the abundance gradually increased one week after the operation and these changes persisted even 6 months after the operation (Paganelli et al., 2019). These changes most likely reflect the changed anatomy of gastrointestinal tract and decreased exposure to gastric acids after operation. Postoperative microbiota changes observed here might reflect the microbiota adaptation to changing conditions in the gastrointestinal tract and energy restriction as well as infeasibilities of fiber-rich foods after bariatric surgery as shown by the increase in Bacteroidetes/Firmicutes ratio and by the reduced abundance of butyrate-producing bacteria. Moreover, a diet rich in resistant starch may not be tolerated after the bariatric surgery. Indeed, a reduction of one of the colonic starch degrading *Ruminococcus obeum*-related bacteria was observed in post-operative microbiota. Based on studies with dietary interventions for weight loss, restrictive diets seem to promote a reduction in intestinal microbiota diversity and a reduction in certain bacterial groups, such as butyrate-producing

Firmicutes, *Clostridium* spp. *Lactobacillus* spp. and *Bifidobacterium* spp. (Damms-Machado et al., 2015; Graessler et al., 2014; Palleja et al., 2016). These changes seem to correlate to a reduction in the intake of complex carbohydrates rather than to a weight loss as such.

Given the fact that bariatric surgery rearranges the gut microbiota, it is worth noting the role of potential modulators of intestinal microbiota, such as prebiotics and synbiotics. In the study of Fernandes's research group oral administration of a prebiotic (fructo-oligosaccharide) after RYGB promoted significant reduction in body weight, whereas both prebiotic and synbiotic supplements were not sufficient to promote significant improvement in the inflammatory markers (Fernandes et al., 2016). Unfortunately, this study lacks the data of change in fecal microbiota. Since bariatric surgery has shown to induce intestinal bacterial overgrowth, Woodard et al. randomized 44 patients undergoing RYGB to placebo or 2.4 billion colonies of *Lactobasillus*. This probiotic administration improved bacterial overgrowth, vitamin B12 availability, and weight loss after RNYGB (Woodard et al., 2009). Based on these findings further investigation in pre- and synbiotics with larger sample size and fecal microbiota sampling is warranted in the future.

During this study, we observed two distinct groups of control subjects with a distinct microbiota composition as well as differences in body-mass index and other host parameters. These groups were not included in the original study design, and hence the findings relating to the distinct control groups in this study are exploratory and limited in sample size. This observation highlights the microbiota signatures as an essential host feature that should be considered already in the study design. These findings should be further verified in an independent study.

There are some limitations in this study. First, study groups were small and while the findings are in line with current literature further research with larger cohorts is warranted. Second, only female subjects were studied, and thus the findings may not be generalized to males. Third, fecal samples were collected in surgery group only before the low-calorie diet and 6 months after the operation. Therefore, we can't rule out that part of the microbiota changes was induced by presurgery diet. Fourth, HITChip is based on cross-hybridization of the complementary 16S rRNA sequences, and may therefore include noise from related sequences, and potential misclassifications between closely related species. For this reason, we have restricted the analysis on genus-level profiles. The other shortcoming of phylogenetic microarrays is that they quantify only the known community members that have been included on the microarray design. The HITChip has been designed to capture all major bacterial groups that have been reported in human gut microbiome; it may miss some rare groups that have low abundance and low population prevalence. On the other hand, this platform is standardized, highly reproducible, and cost efficient compared to sequencing depths that can reach the same sensitivity.

CONCLUSION

The intestinal microbiota composition of morbidly obese subjects shifted towards group A ("normal weight") controls within six months after the bariatric surgery. This shift was detectable in the overall community composition, richness, Bacteroidetes/Firmicutes ratio,

as well as in abundances of specific bacterial genera. However, the individuality of the microbiota community composition was preserved in most cases so that the microbiota profiles could be used to identify individuals before and after surgery. In addition, we observed that the control subjects clustered in two broad groups with a distinct microbiota composition characterized by differences in particular in *Bacteroides*, whose abundance was on average higher in the group A controls than in group B subjects (“lean” controls). Hence, broad community types might be an essential feature to include in the original study design in future studies.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This study was conducted within the Center of Excellence in Cardiovascular and Metabolic Diseases, supported by the Academy of Finland, the University of Turku, the Turku University Hospital, and Åbo Akademi University. This study was supported by the Netherlands Organization for Scientific Research, via the 2008 Spinoza Award and the SIAM Gravitation Grant 024.002.002 to WMdV. Leo Lahti was supported by Academy of Finland (grants 295741, 307127). There was no additional external funding received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
 Academy of Finland, the University of Turku, the Turku University Hospital.
 Åbo Akademi University.
 2008 Spinoza Award.
 SIAM Gravitation Grant: 024.002.002.
 Academy of Finland: 295741, 307127.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Jukka Koffert, Leo Lahti and Lotta Nylund conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Seppo Salminen conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Jarna C. Hannukainen, Paulina Salminen and Pirjo Nuutila conceived and designed the experiments, performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Willem M. de Vos analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.

Human Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

The Ethics Committee of the Hospital District of Southwest Finland approved this research (ETMK 99/180/2010).

Clinical Trial Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

Ethics Committee of the Hospital District of Southwest Finland

Data Availability

The following information was supplied regarding data availability:

Raw data are available in the [Supplementary Files](#).

Clinical Trial Registration

The following information was supplied regarding Clinical Trial registration:

NCT01373892

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.10442#supplemental-information>.

REFERENCES

- Adams TD, Davidson LE, Litwin SE, Kim J, Kolotkin RL, Nanjee MN, Gutierrez JM, Frogley SJ, Ibele AR, Brinton EA, Hopkins PN, McKinlay R, Simper SC, Hunt SC. 2017. Weight and metabolic outcomes 12 years after gastric bypass. *New England Journal of Medicine* 377:1143–1155 DOI 10.1056/NEJMoa1700459.
- Aitchison J. 1986. *The statistical analysis of compositional data, monographs on statistics and applied probability*. London: Chapman & Hall Ltd.
- Allin KH, Tremaroli V, Caesar R, Jensen BAH, Damgaard MTF, Bahl MI, Licht TR, Hansen TH, Nielsen T, Dantoft TM, Linneberg A, Jorgensen T, Vestergaard H, Kristiansen K, Franks PW, consortium I-D, Hansen T, Backhed F, Pedersen O. 2018. Aberrant intestinal microbiota in individuals with prediabetes. *Diabetologia* 61:810–820 DOI 10.1007/s00125-018-4550-1.
- Claesson MJ, O’Sullivan O, Wang Q, Nikkila J, Marchesi JR, Smidt H, De Vos WM, Ross RP, O’Toole PW. 2009. Comparative analysis of pyrosequencing and a phylogenetic microarray for exploring microbial community structures in the human distal intestine. *PLOS ONE* 4:e6669 DOI 10.1371/journal.pone.0006669.
- Collaborators GBDO, Afshin A, Forouzanfar MH, Reitsma MB, Sur P, Estep K, Lee A, Marczak L, Mokdad AH, Moradi-Lakeh M, Naghavi M, Salama JS, Vos T, Abate KH, Abbafati C, Ahmed MB, Al-Aly Z, Alkerwi A, Al-Raddadi R, Amare AT, Amberbir A, Amegah AK, Amini E, Amrock SM, Anjana RM, Arnlov J, Asayesh

- H, Banerjee A, Barac A, Baye E, Bennett DA, Beyene AS, Biadgilign S, Biryukov S, Bjertness E, Boneya DJ, Campos-Nonato I, Carrero JJ, Cecilio P, Cercy K, Ciobanu LG, Cornaby L, Damtew SA, Dandona L, Dandona R, Dharmaratne SD, Duncan BB, Eshрати B, Esteghamati A, Feigin VL, Fernandes JC, Furst T, Gebrehiwot TT, Gold A, Gona PN, Goto A, Habtewold TD, Hadush KT, Hafezi-Nejad N, Hay SI, Horino M, Islami F, Kamal R, Kasaeian A, Katikireddi SV, Kengne AP, Kesavachandran CN, Khader YS, Khang YH, Khubchandani J, Kim D, Kim YJ, Kinfu Y, Kosen S, Ku T, Defo BK, Kumar GA, Larson HJ, Leinsalu M, Liang X, Lim SS, Liu P, Lopez AD, Lozano R, Majeed A, Malekzadeh R, Malta DC, Mazidi M, McAlinden C, McGarvey ST, Mengistu DT, Mensah GA, Mensink GBM, Mezgebe HB, Mirrakhimov EM, Mueller UO, Noubiap JJ, Obermeyer CM, Ogbo FA, Owolabi MO, Patton GC, Pourmalek F, Qorbani M, Rafay A, Rai RK, Ranabhat CL, Reinig N, Safiri S, Salomon JA, Sanabria JR, Santos IS, Sartorius B, Sawhney M, Schmidhuber J, Schutte AE, Schmidt MI, Sepanlou SG, Shamsizadeh M, Sheikhbahaei S, Shin MJ, Shiri R, Shiue I, Roba HS, Silva DAS, Silverberg JL, Singh JA, Stranges S, Swaminathan S, Tabares-Seisdedos R, Tadese F, Tedla BA, Tegegne BS, Terkawi AS, Thakur JS, Tonelli M, Topor-Madry R, Tyrovolas S, Ukwaja KN, Uthman OA, Vaezghasemi M, Vasankari T, Vlassov VV, Vollset SE, Weiderpass E, Werdecker A, Wesana J, Westerman R, Yano Y, Yonemoto N, Yonga G, Zaidi Z, Zenebe ZM, Zipkin B, Murray CJL. 2017. Health effects of overweight and obesity in 195 countries over 25 years. *New England Journal of Medicine* 377:13–27 DOI 10.1056/NEJMoa1614362.
- Cotillard A, Kennedy SP, Kong LC, Prifti E, Pons N, Le Chatelier E, Almeida M, Quinquis B, Levenez F, Galleron N, Gougis S, Rizkalla S, Batto JM, Renault P, consortium ANRM, Dore J, Zucker JD, Clement K, Ehrlich SD. 2013. Dietary intervention impact on gut microbial gene richness. *Nature* 500:585–588 DOI 10.1038/nature12480.
- Dadson P, Ferrannini E, Landini L, Hannukainen JC, Kalliokoski KK, Vaittinen M, Honka H, Karlsson HK, Tuulari JJ, Soinio M, Salminen P, Parkkola R, Pihlajamäki J, Iozzo P, Nuutila P. 2017. Fatty acid uptake and blood flow in adipose tissue compartments of morbidly obese subjects with or without type 2 diabetes: effects of bariatric surgery. *American Journal of Physiology, Endocrinology and Metabolism* 313:E175–E182 DOI 10.1152/ajpendo.00044.2017.
- Damms-Machado A, Mitra S, Schollenberger AE, Kramer KM, Meile T, Konigsrainer A, Huson DH, Bischoff SC. 2015. Effects of surgical and dietary weight loss therapy for obesity on gut microbiota composition and nutrient absorption. *BioMed Research International* 2015:806248 DOI 10.1155/2015/806248.
- Durack J, Lynch SV. 2019. The gut microbiome: relationships with disease and opportunities for therapy. *Journal of Experimental Medicine* 216:20–40 DOI 10.1084/jem.20180448.
- Fernandes R, Beserra BT, Mocellin MC, Kuntz MG, da Rosa JS, de Miranda RC, Schreiber CS, Frode TS, Nunes EA, Trindade EB. 2016. Effects of prebiotic

and synbiotic supplementation on inflammatory markers and anthropometric indices after Roux-en-Y gastric bypass: a randomized, triple-blind, placebo-controlled pilot study. *Journal of Clinical Gastroenterology* 50:208–217 DOI [10.1097/MCG.0000000000000328](https://doi.org/10.1097/MCG.0000000000000328).

- Forslund K, Hildebrand F, Nielsen T, Falony G, Le Chatelier E, Sunagawa S, Prifti E, Vieira-Silva S, Gudmundsdottir V, Pedersen HK, Arumugam M, Kristiansen K, Voigt AY, Vestergaard H, Hercog R, Costea PI, Kultima JR, Li J, Jorgensen T, Levenez F, Dore J, Meta HITC, Nielsen HB, Brunak S, Raes J, Hansen T, Wang J, Ehrlich SD, Bork P, Pedersen O. 2015. Disentangling type 2 diabetes and metformin treatment signatures in the human gut microbiota. *Nature* 528:262–266 DOI [10.1038/nature15766](https://doi.org/10.1038/nature15766).
- Furet JP, Kong LC, Tap J, Poitou C, Basdevant A, Bouillot JL, Mariat D, Corthier G, Dore J, Henegar C, Rizkalla S, Clement K. 2010. Differential adaptation of human gut microbiota to bariatric surgery-induced weight loss: links with metabolic and low-grade inflammation markers. *Diabetes* 59:3049–3057 DOI [10.2337/db10-0253](https://doi.org/10.2337/db10-0253).
- Graessler J, Bornstein TD, Goel D, Bhalla VP, Lohmann T, Wolf T, Koch M, Qin Y, Licinio J, Wong ML, Chavakis T, Xu A, Shevchenko A, Schuhmann K, Schwarz PE, Schulte KM, Patel A, Bornstein SR. 2014. Lipidomic profiling before and after Roux-en-Y gastric bypass in obese patients with diabetes. *Pharmacogenomics Journal* 14:201–207 DOI [10.1038/tpj.2013.42](https://doi.org/10.1038/tpj.2013.42).
- Healey GR, Murphy R, Brough L, Butts CA, Coad J. 2017. Interindividual variability in gut microbiota and host response to dietary interventions. *Nutrition Reviews* 75:1059–1080 DOI [10.1093/nutrit/nux062](https://doi.org/10.1093/nutrit/nux062).
- Ilhan ZE, DiBaise JK, Isern NG, Hoyt DW, Marcus AK, Kang DW, Crowell MD, Rittmann BE, Krajmalnik-Brown R. 2017. Distinctive microbiomes and metabolites linked with weight loss after gastric bypass, but not gastric banding. *ISME Journal* 11:2047–2058 DOI [10.1038/ismej.2017.71](https://doi.org/10.1038/ismej.2017.71).
- Karlsson FH, Tremaroli V, Nookaew I, Bergstrom G, Behre CJ, Fagerberg B, Nielsen J, Backhed F. 2013. Gut metagenome in European women with normal, impaired and diabetic glucose control. *Nature* 498:99–103 DOI [10.1038/nature12198](https://doi.org/10.1038/nature12198).
- Karmi A, Iozzo P, Viljanen A, Hirvonen J, Fielding BA, Virtanen K, Oikonen V, Kemppainen J, Viljanen T, Guiducci L, Haaparanta-Solin M, Nagren K, Solin O, Nuutila P. 2010. Increased brain fatty acid uptake in metabolic syndrome. *Diabetes* 59:2171–2177 DOI [10.2337/db09-0138](https://doi.org/10.2337/db09-0138).
- Koffert J, Stahle M, Karlsson H, Iozzo P, Salminen P, Roivainen A, Nuutila P. 2018. Morbid obesity and type 2 diabetes alter intestinal fatty acid uptake and blood flow. *Diabetes, Obesity and Metabolism* 20:1384–1390 DOI [10.1111/dom.13228](https://doi.org/10.1111/dom.13228).
- Kolmeder CA, Ritari J, Verdam FJ, Muth T, Keskitalo S, Varjosalo M, Fuentes S, Greve JW, Buurman WA, Reichl U, Rapp E, Martens L, Palva A, Salonen A, Rensen SS, De Vos WM. 2015. Colonic metaproteomic signatures of active bacteria and the host in obesity. *Proteomics* 15:3544–3552 DOI [10.1002/pmic.201500049](https://doi.org/10.1002/pmic.201500049).
- Kootte RS, Vrieze A, Holleman F, Dallinga-Thie GM, Zoetendal EG, De Vos WM, Groen AK, Hoekstra JB, Strees ES, Nieuwdorp M. 2012. The therapeutic potential

- of manipulating gut microbiota in obesity and type 2 diabetes mellitus. *Diabetes Obes Metab* 14:112–120 DOI 10.1111/j.1463-1326.2011.01483.x.
- Krajmalnik-Brown R, Ilhan ZE, Kang DW, DiBaise JK. 2012.** Effects of gut microbes on nutrient absorption and energy regulation. *Nutrition in Clinical Practice* 27:201–214 DOI 10.1177/0884533611436116.
- Laferrere B, Teixeira J, McGinty J, Tran H, Egger JR, Colarusso A, Kovack B, Bawa B, Koshy N, Lee H, Yapp K, Olivan B. 2008.** Effect of weight loss by gastric bypass surgery versus hypocaloric diet on glucose and incretin levels in patients with type 2 diabetes. *Journal of Clinical Endocrinology and Metabolism* 93:2479–2485 DOI 10.1210/jc.2007-2851.
- Lahti L, Elo LL, Aittokallio T, Kaski S. 2011.** Probabilistic analysis of probe reliability in differential gene expression studies with short oligonucleotide arrays. *IEEE/ACM Trans Comput Biol Bioinform* 8:217–225 DOI 10.1109/TCBB.2009.38.
- Lahti L, Salojärvi J, Salonen A, Scheffer M, De Vos WM. 2014.** Tipping elements in the human intestinal ecosystem. *Nature Communications* 5:4344 DOI 10.1038/ncomms5344.
- Lahti L, Torrente A, Elo LL, Brazma A, Rung J. 2013.** A fully scalable online pre-processing algorithm for short oligonucleotide microarray atlases. *Nucleic Acids Research* 41:e110 DOI 10.1093/nar/gkt229.
- Lee CJ, Florea L, Sears CL, Maruthur N, Potter JJ, Schweitzer M, Magnuson T, Clark JM. 2019.** Changes in gut microbiome after bariatric surgery versus medical weight loss in a pilot randomized trial. *Obesity Surgery* 29:3239–3245 DOI 10.1007/s11695-019-03976-4.
- Ley RE, Turnbaugh PJ, Klein S, Gordon JI. 2006.** Microbial ecology: human gut microbes associated with obesity. *Nature* 444:1022–1023 DOI 10.1038/4441022a.
- Liou AP, Paziuk M, Luevano Jr JM, Machineni S, Turnbaugh PJ, Kaplan LM. 2013.** Conserved shifts in the gut microbiota due to gastric bypass reduce host weight and adiposity. *Science Translational Medicine* 5:178ra141 DOI 10.1126/scitranslmed.3005687.
- Louis BP, Maron PA, Menasseri-Aubry S, Sarr A, Leveque J, Mathieu O, Jolivet C, Leterme P, Viaud V. 2016.** Microbial diversity indexes can explain soil carbon dynamics as a function of carbon source. *PLOS ONE* 11:e0161251 DOI 10.1371/journal.pone.0161251.
- McMurdie PJ, Holmes S. 2013.** phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLOS ONE* 8:e61217 DOI 10.1371/journal.pone.0061217.
- Nylund L, Heilig HG, Salminen S, De Vos WM, Satokari R. 2010.** Semi-automated extraction of microbial DNA from feces for qPCR and phylogenetic microarray analysis. *Journal of Microbiological Methods* 83:231–235 DOI 10.1016/j.mimet.2010.09.003.
- Paganelli FL, Luyer M, Hazelbag CM, Uh HW, Rogers MRC, Adriaans D, Berbers RM, Hendrickx APA, Viveen MC, Groot JA, Bonten MJM, Fluit AC, Willems RJL, Leavis HL. 2019.** Roux-Y gastric bypass and sleeve gastrectomy directly change gut

- microbiota composition independent of surgery type. *Scientific Reports* 9:10979 DOI 10.1038/s41598-019-47332-z.
- Palleja A, Kashani A, Allin KH, Nielsen T, Zhang C, Li Y, Brach T, Liang S, Feng Q, Jorgensen NB, Bojsen-Moller KN, Dirksen C, Burgdorf KS, Holst JJ, Madsbad S, Wang J, Pedersen O, Hansen T, Arumugam M. 2016. Roux-en-Y gastric bypass surgery of morbidly obese patients induces swift and persistent changes of the individual gut microbiota. *Genome Medicine* 8:67 DOI 10.1186/s13073-016-0312-1.
- Patil DP, Dhotre DP, Chavan SG, Sultan A, Jain DS, Lanjekar VB, Gangawani J, Shah PS, Todkar JS, Shah S, Ranade DR, Patole MS, Shouche YS. 2012. Molecular analysis of gut microbiota in obesity among Indian individuals. *Journal of Biosciences* 37:647–657 DOI 10.1007/s12038-012-9244-0.
- Pedersen HK, Gudmundsdottir V, Nielsen HB, Hyotylainen T, Nielsen T, Jensen BA, Forslund K, Hildebrand F, Prifti E, Falony G, Le Chatelier E, Levenez F, Dore J, Mattila I, Plichta DR, Poho P, Hellgren LI, Arumugam M, Sunagawa S, Vieira-Silva S, Jorgensen T, Holm JB, Trost K, Meta HITC, Kristiansen K, Brix S, Raes J, Wang J, Hansen T, Bork P, Brunak S, Oresic M, Ehrlich SD, Pedersen O. 2016. Human gut microbes impact host serum metabolome and insulin sensitivity. *Nature* 535:376–381 DOI 10.1038/nature18646.
- Plum L, Ahmed L, Febres G, Bessler M, Inabnet W, Kunreuther E, McMahon DJ, Korner J. 2011. Comparison of glucostatic parameters after hypocaloric diet or bariatric surgery and equivalent weight loss. *Obesity (Silver Spring)* 19:2149–2157 DOI 10.1038/oby.2011.134.
- Puzziferri N, Roshek 3rd TB, Mayo HG, Gallagher R, Belle SH, Livingston EH. 2014. Long-term follow-up after bariatric surgery: a systematic review. *Journal of the American Medical Association* 312:934–942 DOI 10.1001/jama.2014.10706.
- Qin J, Li Y, Cai Z, Li S, Zhu J, Zhang F, Liang S, Zhang W, Guan Y, Shen D, Peng Y, Zhang D, Jie Z, Wu W, Qin Y, Xue W, Li J, Han L, Lu D, Wu P, Dai Y, Sun X, Li Z, Tang A, Zhong S, Li X, Chen W, Xu R, Wang M, Feng Q, Gong M, Yu J, Zhang Y, Zhang M, Hansen T, Sanchez G, Raes J, Falony G, Okuda S, Almeida M, Le Chatelier E, Renault P, Pons N, Batto JM, Zhang Z, Chen H, Yang R, Zheng W, Li S, Yang H, Wang J, Ehrlich SD, Nielsen R, Pedersen O, Kristiansen K, Wang J. 2012. A metagenome-wide association study of gut microbiota in type 2 diabetes. *Nature* 490:55–60 DOI 10.1038/nature11450.
- Rajilic-Stojanovic M, Heilig HG, Molenaar D, Kajander K, Surakka A, Smidt H, De Vos WM. 2009. Development and application of the human intestinal tract chip, a phylogenetic microarray: analysis of universally conserved phylotypes in the abundant microbiota of young and elderly adults. *Environmental Microbiology* 11:1736–1751 DOI 10.1111/j.1462-2920.2009.01900.x.
- Remely M, Hippe B, Zanner J, Aumüller E, Brath H, Haslberger AG. 2016. Gut microbiota of obese, type 2 diabetic individuals is enriched in faecalibacterium prausnitzii, akkermansia muciniphila and peptostreptococcus anaerobius after weight loss. *Endocrine, Metabolic & Immune Disorders - Drug Targets* 16:99–106 DOI 10.2174/1871530316666160831093813.

- Ridaura VK, Faith JJ, Rey FE, Cheng J, Duncan AE, Kau AL, Griffin NW, Lombard V, Henrissat B, Bain JR, Muehlbauer MJ, Ilkayeva O, Semenkovich CF, Funai K, Hayashi DK, Lyle BJ, Martini MC, Ursell LK, Clemente JC, Van Treuren W, Walters WA, Knight R, Newgard CB, Heath AC, Gordon JI. 2013. Gut microbiota from twins discordant for obesity modulate metabolism in mice. *Science* 341:1241214 DOI 10.1126/science.1241214.
- Sanchez-Alcoholado L, Gutierrez-Repiso C, Gomez-Perez AM, Garcia-Fuentes E, Tinahones FJ, Moreno-Indias I. 2019. Gut microbiota adaptation after weight loss by Roux-en-Y gastric bypass or sleeve gastrectomy bariatric surgeries. *Surgery for Obesity and Related Diseases* 15:1888–1895 DOI 10.1016/j.soard.2019.08.551.
- Schauer PR, Bhatt DL, Kirwan JP, Wolski K, Aminian A, Brethauer SA, Navaneethan SD, Singh RP, Pothier CE, Nissen SE, Kashyap SR, Investigators S. 2017. Bariatric surgery versus intensive medical therapy for diabetes - 5-year outcomes. *New England Journal of Medicine* 376:641–651 DOI 10.1056/NEJMoa1600869.
- Sjostrom L. 2013. Review of the key results from the Swedish Obese Subjects (SOS) trial - a prospective controlled intervention study of bariatric surgery. *Journal of Internal Medicine* 273:219–234 DOI 10.1111/joim.12012.
- Turnbaugh PJ, Hamady M, Yatsunenko T, Cantarel BL, Duncan A, Ley RE, Sogin ML, Jones WJ, Roe BA, Affourtit JP, Egholm M, Henrissat B, Heath AC, Knight R, Gordon JI. 2009. A core gut microbiome in obese and lean twins. *Nature* 457:480–484 DOI 10.1038/nature07540.
- Viljanen AP, Iozzo P, Borra R, Kankaanpaa M, Karmi A, Lautamaki R, Jarvisalo M, Parkkola R, Ronnema T, Guiducci L, Lehtimaki T, Raitakari OT, Mari A, Nuutila P. 2009. Effect of weight loss on liver free fatty acid uptake and hepatic insulin resistance. *Journal of Clinical Endocrinology and Metabolism* 94:50–55 DOI 10.1210/jc.2008-1689.
- Woodard GA, Encarnacion B, Downey JR, Peraza J, Chong K, Hernandez-Boussard T, Morton JM. 2009. Probiotics improve outcomes after Roux-en-Y gastric bypass surgery: a prospective randomized trial. *Journal of Gastrointestinal Surgery* 13:1198–1204 DOI 10.1007/s11605-009-0891-x.